

The role of CCR1 and therapeutic effects of anti-CCL3 antibody in herpes simplex virus-induced Behçet's disease mouse model

Hasan M. Sayeed,¹  Eun-So Lee MD, PhD,² Hae-Ok Byun PhD³ and Seonghyang Sohn PhD^{1,3}

¹Department of Biomedical Sciences, Ajou University School of Medicine, Suwon, ,

²Department of Dermatology, Ajou University School of Medicine, Suwon and ³Department of Microbiology, Ajou University School of Medicine, Suwon, Korea

doi:10.1111/imm.13102

Received 22 November 2018; revised 12 June 2019; accepted 25 June 2019.

Correspondence: Prof. Seonghyang Sohn, Yeongtong-gu, Worldcup-ro, 164, Suwon 16499, Korea. Email: sohnsh@ajou.ac.kr
Senior author: Seonghyang Sohn

Introduction

Behçet's disease (BD) is a chronic systemic inflammatory disease with unclear etiopathogenesis. Gene variants of CC chemokine receptor type 1 (CCR1) in patients with BD have been identified by genome-wide association studies.^{1,2} However, the CCR1 protein expression in patients with BD or in mouse models remains unclear.

CC chemokine receptor type 1 is expressed in cells of the immune system, including lymphocytes, monocytes, neutrophils and dendritic cells.^{3,4} It is involved in host defense, hematopoiesis, T helper type 1/2 (Th1/2) cell balance, and chemotaxis.^{5,6} The CCR1 expression is correlated with various inflammatory diseases, including

Summary

Behçet's disease (BD) is a chronic systemic inflammatory disease with unclear etiopathogenesis. Although gene variants of CC chemokine receptor type 1 (CCR1) have been reported, the protein expression of CCR1 in patients with BD remains unclear. The objective of this study was to analyze the frequencies of CCR1⁺ cells in a herpes simplex virus-induced mouse model of BD. The frequencies of CCR1⁺ cells on the surface and in the cytoplasm of peripheral blood mononuclear cells and lymph nodes were analyzed by flow cytometry. The CCR1⁺ cells were significantly down-regulated in BD mice compared with the normal control and symptom-free control mice. Colchicine and pentoxifylline treatment improved the symptoms of BD and increased the frequencies of CCR1⁺ cells in BD mice. Treatment with chemokine CC motif ligand 3 (CCL3), a ligand of CCR1, caused BD symptoms to deteriorate in 10 of 16 BD mice (62.5%) via down-regulation of CCR1⁺ cells. Anti-CCL3 antibody treatment ameliorated BD symptoms in 10 of 20 mice (50%) and significantly decreased the disease severity score compared with CCL3-treated BD mice ($P = 0.01$) via up-regulation of CCR1⁺ cell frequencies. In patients with BD, plasma levels of CCL3 in an active state were significantly higher than in healthy control individuals ($P = 0.02$). These results show that the up-regulation of CCR1⁺ cells was related to the control of systemic inflammation of BD in mouse models.

Keywords: Behçet's disease; CCL3; CCR1; herpes simplex virus; inflammation; mouse model.

rheumatoid arthritis,⁷ asthma⁸ and lupus nephritis.⁹ CCR1 knockout (−/−) mice show reduced pulmonary inflammation,¹⁰ and prolonged allograft survival.¹¹ CCR1 polymorphism might be a general risk factor for autoimmune or inflammatory disease. CCR1 antagonists have been used in clinical trials to treat multiple sclerosis, psoriasis and rheumatoid arthritis.¹²

Herpes simplex virus (HSV) has been reported as an etiological factor in BD.^{13–15} A mouse model of HSV-induced inflammatory BD was developed and used in various studies.^{16–19} The objective of the present study was to determine the expression of CCR1 in mice with inflammatory BD symptoms and to investigate whether CCR1 was regulated by ligands, agonists or antagonists. Flow

Abbreviation: BD, Behçet's disease; BDN, herpes simplex virus-inoculated, but asymptomatic healthy mice; CCL3, chemokine CC motif ligand 3; CCR1, CC chemokine receptor type 1; GM-CSF, granulocyte–macrophage colony-stimulating factor; IL-10, interleukin-10

cytometry was used to examine the frequencies of CCR1⁺ cells on the surface and in the cytoplasm of lymph node (LN) cells and peripheral blood mononuclear cells (PBMC). Our results demonstrated the correlation between CCR1 surface expression on PBMC and the severity of BD symptoms in mice.

Materials and methods

Animal experiments

Male ICR mice (4 weeks old) were infected with HSV type 1 (1×10^6 plaque-forming units/ml, F strain) grown in Vero cells as described previously.¹⁶ The virus was inoculated twice into mice at a frequency of 10 days. After the second HSV inoculation, mice were observed from week 4 to week 32. Animals were handled in accordance with guidelines provided by the Institutional Animal Care and Use Committee of Ajou University School of Medicine (AMC-2016-0010).

Symptoms of BD in mice

Among all the mice infected with HSV, 15% developed BD-like symptoms including skin ulcerations, skin pustules, genital ulcerations, erythema, oral ulceration, ear inflammation, arthritis, genital ulcerations, red eye (left, right), reduced vision (left, right), weight loss and facial swelling. Oral, genital and skin ulcers, and eye symptoms were classified as major symptoms. Other symptoms were identified as minor symptoms. Mice displaying more than one major and one minor symptom were diagnosed with BD. Each symptom score was assigned a single point. The sum of all symptom scores was used to determine the severity of BD. Disappearance of symptoms or $\geq 20\%$ decrease in lesion size was classified as symptom improvement. The disease severity score of BD was calculated according to the BD activity index outlined in the BD Activity Form (www.behcet.ws/pdf/BehcetsDiseaseActivityForm.pdf). HSV-inoculated, but asymptomatic, healthy mice (BDN mice) served as the control group, as described previously.

Flow cytometry

Peripheral blood mononuclear cells and LN cells were isolated from mice. Erythrocytes were eliminated with ammonium-chloride-potassium solution and washed with phosphate-buffered saline (PBS) followed by surface staining of 1×10^6 cells with mouse CCR1 allophycocyanin-conjugated antibody (cat# FAB5986A; R&D Systems, Minneapolis, MN) for 30 min at 4° in the dark. For intracytoplasmic staining, cells were fixed in intracytoplasmic fixation buffer for 20 min at room temperature. After washing twice with a permeabilization buffer, 1×10^6 cells were stained with an anti-mouse CCR1 antibody,

followed by analysis of these cells with a flow cytometer (FACS Canto II; Becton Dickinson, San Jose, CA).

Western blot analysis

Fresh PBMC were washed twice with cold PBS and lysed with lysis buffer (50 mM Tris-Cl, pH 7.4, 0.15 M NaCl, 1 mM EDTA, 1% Triton X-100, 10 µg/ml each of aprotinin and leupeptin, and 1 mM phenylmethylsulphonyl fluoride). The protein (30 µg) lysate was electrophoresed on a 10% sodium dodecyl sulfate-polyacrylamide gel. The proteins were transferred onto nitrocellulose membranes (Protran; GE Healthcare Life Sciences, Chicago, IL, USA) and subjected to immunoblot analysis using antibodies. The immunoblots were developed with the enhanced chemiluminescence system (GE Healthcare Life Sciences). Antibody against CCR1 (MAB5986) was purchased from R&D Systems and glyceraldehyde 3-phosphate dehydrogenase (GAPDH) antibody (sc-32233) was from Santa Cruz Biotechnology, Inc. (Dallas, TX).

Administration of cytokines, ligand and antagonist

Normal and BD mice were treated *in vivo* with PBS, interleukin-10 (IL-10) (50 pg/mouse/day) (cat# CYT-497; Prospecbio, Ness-Ziona, Israel), granulocyte-macrophage colony-stimulating factor (GM-CSF) (200 pg/mouse/day) (cat# CYT-222; Prospecbio), chemokine CC motif ligand 3 (CCL3) (100 pg/mouse/day) (cat# CHM-320; Prospecbio), or anti-CCL3 antibody (20 ng/mouse/day) (cat# AF-450-NA; R&D Systems) for five consecutive days by intraperitoneal administration. Normal mice were 3.5 weeks old. PBMC and LN cells were isolated 3 hr after the last medication for analysis of CCR1⁺ cells by flow cytometry. BX471, a CCR1 antagonist (cat# SML0020; Sigma-Aldrich, St Louis, MO), was intraperitoneally administered to normal and BD mice at 0.5 mg/mouse three times (every other day). BD symptoms were observed before and after treatment. Images were obtained to document the treatment-related changes.

Drug treatments

Mice were treated orally with PBS, colchicine (2 µg/mouse/day), or pentoxifylline (400 µg/mouse/day) for five consecutive days. Changes in symptoms were monitored. At 3 hr after the last medication, the mice were killed by intramuscular injection into the hind leg with a ketamine/xylazine cocktail (15 mg/kg ketamine and 10 mg/kg xylazine). PBMC and LN cells were isolated for analysis of CCR1⁺ cells.

Patients with BD

The patient population consisted of 12 patients with BD, who presented for the first time or were monitored at the

Department of Dermatology, Ajou University Hospital. Sampling from patients with BD was random when the patients first arrived. Clinical characteristics and therapeutic histories of these patients are shown in Tables 1 and 2. According to the International Study Group for BD criteria, the presence of any two of the following symptoms, in addition to recurrent oral ulcerations, is considered as BD diagnosis: recurrent genital ulceration, cutaneous erythema nodosum, large-vessel vasculitis, arthritis, uveitis, and/or a positive pathergy test. The disease severity score was calculated from Behçet's disease current activity form 2006 (<http://www.behcetdiseasesociety.org/behcetwsData/Uploads/files/BehcetsDiseaseActivityForm.pdf>). Patients with active BD ($n = 12$, 11 men, one woman, 45.0 ± 10.2 years) were enrolled. Gender differences among BD patients with mucocutaneous manifestations were inconsistent among different nationalities.²⁰ Informed consent was obtained from patients before the study. The healthy control group ($n = 12$, 48.8 ± 9.2 years) comprised one man and 11 women. Serum was provided by the Biobank of Ajou University Hospital, a member of the Korea Biobank Network. This study was approved by the ethics committee of Ajou University Medical Center Institutional Review Board (AJIRB-GN3-07-098) and all experiments were conducted according to the Institutional Review Board permitted protocol.

CCL3 ELISA

Peripheral blood was collected in EDTA-coated test tubes and centrifuged at 1000 g for 5 min at 20°. The

supernatant was stored at -70° until used. CCL3 levels were measured in plasma using a commercial enzyme-linked immunosorbent assay (ELISA) kit (Cat# BMS2029INST; Thermo Fisher Scientific, Waltham, MA) following the manufacturer's instructions. For ELISA analysis, each sample was applied in duplicate wells and three times in different ELISA plates.

Statistical analysis

The data and statistical analysis complied with the recommendations of experimental design and analysis in pharmacology.

Results

Frequencies of CCR1⁺ cells in BD mice

Flow cytometric analysis was performed to analyze the surface and cytoplasm CCR1⁺ cells stained with anti-CCR1 antibody for PBMC and LN cells isolated from normal ($n = 10$), BDN ($n = 10$), and BD ($n = 10$) mice. Frequencies of CCR1⁺ cells on the surface of PBMC were significantly down-regulated in BD mice compared with those in normal mice ($7.06 \pm 1.8\%$ versus $11.11 \pm 2.5\%$, $P = 0.0005$) or BDN mice ($7.06 \pm 1.8\%$ versus $10.58 \pm 1.92\%$, $P = 0.0006$). Frequencies of CCR1⁺ cells on the surface of LN cells were also down-regulated in BD mice compared with normal mice ($10.6 \pm 3.2\%$ versus $12.99 \pm 1.5\%$, $P = 0.05$) or BDN mice ($10.6 \pm 3.2\%$ versus $11.12 \pm 2.1\%$, $P = 0.67$) (Fig. 1a).

Table 1. Clinical characteristics of patients with Behçet's disease

Patient no.	Age	Sex	OU	GU	Skin	OL	Arth	Neur	Vas	Path	HLA
Before treatment											
1	46	M	+	+	+	+	+				+
2	43	M	+		+				+		-
3	20	M	+	+	+	+					-
4	36	M	+	+	+	+		+			+
5	60	M	+	+	+	+				-	-
6	44	M	+	+		+					-
7	52	M	+		+					+	
8	45	F	+	+	+	+				-	-
9	52	M	+		+				+	+	+
10	47	M	+	+	+	+				-	+
11	54	M	+		+				+	+	-
12	41	M	+	+		+			+		-
After treatment											
2	43	M	+		+						-
3	20	M	+		+	+					-
6	44	M	+		+						-
7	52	M	+							+	
8	45	F	+							-	-

Arth, Arthritis; GU, genital ulcers; HLA, HLA-B51; Neur, neurological involvement; OL, ocular lesion; OU, oral ulcers; Path, pathergy test; Skin, skin lesion; Vas, vasculitis.

Table 2. Therapeutic history of patients with Behçet's disease

Patient	Colchicine	Azathioprine	Salazopyrine	Pentoxifylline	Minocycline	Aspirin	Vit.D	Methotrexate	Prednisolone	Methylprednisolone
1	+	+		+						
2	+			+	+				+	
3	+	+								
4	+		+	+		+	+			
5	+									
6	+			+	+			+	+	
7	+			+				+		
8	+		+	+				+		
9	+			+			+			
10	+			+						+
11	+			+						
12	+									




Figure 1. Frequencies of CCR1⁺ cells in normal, herpes simplex virus-inoculated, but asymptomatic healthy (BDN) and Behçet's disease (BD) mice. Peripheral blood mononuclear cells (PBMC) and lymph node cells were isolated from normal, BDN and BD mice. Frequencies of surface CCR1⁺ cells (a) and cytoplasm CCR1⁺ cells (b) of whole leukocytes were analyzed by FACS. Representative flow cytometric histograms of surface CCR1⁺ PBMC and lymph node cells are shown (c). (d) Protein expression of CCR1 by Western blot, (e) Expression ratio of CCR1 to GAPDH.

The frequencies of cytoplasm staining in CCR1⁺ PBMC were significantly lower in BD mice than in normal mice (32.08 ± 16.3% versus 73.6 ± 3%, $P < 0.0001$) or BDN

mice (32.08 ± 16.3% versus 58.4 ± 11.05%, $P = 0.0005$). The frequencies of CCR1⁺ LN cells were also significantly lower in BD mice than in normal mice

(58.04 ± 22.39% versus 82.14 ± 8.15%, $P = 0.005$). There were no differences in frequencies of CCR1⁺ cells between BD and BDN mice (58.04 ± 22.39% versus 59.10 ± 27.6, $P = 0.9$) (Fig. 1b). Representative histograms of surface staining results are presented in Fig. 1(c). Figure 1(d) shows the Western blot analysis for CCR1 expression in whole PBMC lysates in normal, BDN and BD mice ($n = 3$ in each group). The expression ratio of CCR1 compared with GAPDH was not different between BDN (1.42 ± 0.34) and BD (1.48 ± 1.15) mice (Fig. 1e, $n = 7$). This means the entire amount of CCR1 in PBMC was not correlated with the symptoms.

Interleukin-10 and GM-CSF affect frequencies of CCR1⁺ cells in normal mice

Interleukin-10 synergistically enhanced GM-CSF-induced CCR1.²¹ GM-CSF is known as a CCR1-enhancing cytokine.²² Therefore, IL-10 and GM-CSF were administered to normal and BD mice. To determine the functional doses in mice *in vivo*, IL-10 and GM-CSF were injected into 3.5-week-old normal mice intraperitoneally at the following doses: IL-10 (25, 50 and 100 pg) and GM-CSF (50, 100 and 200 pg). At 3 hr after the last treatment, PBMC were isolated from mice for flow cytometry to measure the frequencies of CCR1⁺ cells on the surface and cytoplasm.

In mice treated with 50 pg of IL-10 ($n = 13$), the frequencies of surface CCR1⁺ cells were significantly up-regulated in PBMC compared with PBS-treated control mice ($n = 5$) (13.14 ± 0.9% versus 11.0 ± 0.3%, $P = 0.009$). Treatment with 25 pg of IL-10 ($n = 9$) did not affect the frequencies of CCR1⁺ cells compared with PBS-treated control mice ($n = 5$) (11.74 ± 0.96% versus 11.0 ± 0.3%, $P = 0.47$). Treatment with 100 pg of IL-10 ($n = 6$) up-regulated the frequencies of CCR1⁺ cells compared with PBS-treated control mice ($n = 5$), although the difference was not statistically significant (13.65 ± 2.05% versus 11.0 ± 0.3%, $P = 0.18$). The frequencies of cytoplasm CCR1⁺ PBMC were slightly increased in mice treated with 50 pg of IL-10 and 100 pg of IL-10 compared with those in PBS-treated control mice (62.575 ± 9.89% versus 56.05 ± 23.12%, $P = 0.6$; 58.8 ± 23.33% versus 56.05 ± 23.12%, $P = 0.9$), although the differences were not statistically significant (Fig. 2a).

In normal mice exposed to 200 pg of GM-CSF ($n = 5$), the frequencies of surface CCR1⁺ PBMC were up-regulated compared with PBS-treated control mice (15.65 ± 1.4% versus 11.0 ± 0.42%, $P = 0.05$). In mice treated with 100 pg of GM-CSF ($n = 6$), the frequencies of CCR1⁺ cells were higher compared with the levels in PBS-treated control mice ($n = 5$) (13.75 ± 1.34% versus 11.0 ± 0.3%, $P = 0.11$). However, the difference was not statistically significant. Mice treated with 50 pg of GM-

CSF ($n = 6$) showed frequencies of CCR1⁺ cells similar to those in PBS-treated control mice ($n = 5$). In the cytoplasm of PBMC, frequencies of CCR1⁺ cells in mice treated with GM-CSF at 50, 100 and 200 pg (62.25 ± 3.18%; 63.95 ± 1.9%; 64.7 ± 1.5%, respectively) were not significantly ($P > 0.05$) different compared with those in the PBS control group (56.05 ± 23.12%) (Fig. 2b).

Interleukin-10 and GM-CSF up-regulate frequencies of CCR1⁺ cells in BD mice

According to functional doses of IL-10 and GM-CSF in normal mice, 50 pg/mouse/day of IL-10 and 200 pg/mouse/day of GM-CSF were administered to BD mice to up-regulate CCR1⁺ cells. After five consecutive days of treatment to BD mice, PBMC and LN cells were subjected to surface and cytoplasm staining for FACS analysis. In the surface of PBMC, the frequencies of CCR1⁺ cells in BD mice treated with IL-10 ($n = 7$) and GM-CSF ($n = 7$) showed significant up-regulation compared with those in PBS-treated BD mice ($n = 10$) (IL-10 versus PBS: 13.61 ± 2.5% versus 7.06 ± 1.8%, $P < 0.0001$; GM-CSF versus PBS: 9.0 ± 1.3% versus 7.06 ± 1.8%, $P = 0.03$). The surface CCR1⁺ LN cells were significantly up-regulated compared with PBS-treated BD mice (IL-10 versus PBS: 17.6 ± 2.7% versus 10.6 ± 3.23%, $P = 0.0002$; GM-CSF versus PBS: 15.4 ± 1.6% versus 10.6 ± 3.23%, $P = 0.003$) (Fig. 2c). In the cytoplasm, CCR1⁺ PBMC also increased after treatment with IL-10 (54.67 ± 5.00% versus 32.04 ± 16.3%, $P = 0.02$) and GM-CSF (82.1 ± 6.4% versus 32.08 ± 16.3%, $P = 0.001$) compared with PBS-treated BD mice. Cytoplasm CCR1⁺ cells in LN were increased only by GM-CSF (77.85 ± 5.9% versus 58.04%, $P = 0.04$). Frequencies of cytoplasm CCR1⁺ LN cells were not changed after IL-10 treatment (Fig. 2d). IL-10 and GM-CSF up-regulated the frequencies of CCR1⁺ cells in both normal and BD mice.

CCL3 down-regulates the expression of CCR1⁺ cells in normal mice

CCR1 has multiple ligands. CCL3 is one of its ligands for CCR1 down-regulation function.²³ To determine whether CCR1 was regulated by CCL3, CCL3 was intraperitoneally injected into normal mice. Frequencies of CCR1⁺ PBMC and LN cells were decreased after CCL3 treatment. In surface staining of PBMC, treatment with 100 pg of CCL3 ($n = 6$) significantly decreased the frequencies of CCR1⁺ cells compared with PBS control ($n = 5$) (4.62 ± 1.3% versus 12.92 ± 1.89%, $P = 0.0002$). Treatment with 50 pg CCL3 decreased frequencies of CCR1⁺ cells, although the decrease was not significant (9.05 ± 2.7% versus 12.92 ± 1.89%, $P = 0.06$). In surface staining of LN cells, treatment with CCL3 at 100 pg ($n = 6$) significantly down-regulated the frequencies of CCR1⁺ cells




Figure 2. Frequencies of CCR1⁺ cells in cytokine-treated normal and Behçet's disease (BD) mice. Interleukin-10 (IL-10) (25, 50 and 100 pg) and granulocyte-macrophage colony-stimulating factor (GM-CSF) (50, 100 and 200 pg) were intraperitoneally injected into normal mice for five consecutive days and peripheral blood mononuclear cells (PBMC) were isolated. Frequencies of CCR1⁺ cells were analyzed by FACS for expression on the surface and in the cytoplasm of mice after treatment with IL-10 (a) or GM-CSF (b). IL-10 (50 pg) and GM-CSF (200 pg) were intraperitoneally injected into BD mice for five consecutive days. PBMC and lymph node cells were isolated. Frequencies of surface CCR1⁺ (c) and cytoplasm CCR1⁺ (d) PBMC and lymph node cells derived from IL-10- and GM-CSF-treated BD mice were analyzed by FACS.

compared with PBS control ($n = 5$) ($9.42 \pm 1.32\%$ versus $13.15 \pm 1.7\%$, $P = 0.009$). Treatment with CCL3 50 pg ($n = 6$) resulted in similar frequencies of CCR1⁺ cells compared with PBS-treated control mice (Fig. 3a).

The frequencies of cytoplasm CCR1⁺ PBMC after treatment with CCL3 at 50 and 100 pg were significantly decreased to ($50.95 \pm 4.3\%$, $P < 0.0001$) and ($33.9 \pm 4.4\%$, $P < 0.0001$) compared with PBS control ($74.05 \pm 3.4\%$). Cytoplasm CCR1⁺ LN cells were significantly down-regulated in mice treated with 50 pg of CCL3 ($25.65 \pm 1.34\%$, $P < 0.0001$) or 100 pg of CCL3 ($39.48 \pm 2.4\%$, $P < 0.0001$) compared with PBS-treated control mice ($73.75 \pm 1.4\%$) (Fig. 3b). A histogram of surface and cytoplasm CCR1⁺ PBMC after treatment with CCL3 is presented in Fig. 3(c).

CCL3 down-regulates the expression of CCR1⁺ cells and deteriorates symptoms in BD mice

To down-regulate CCR1⁺ cells in BD mice, CCL3 100 pg/day was administered to BD mice ($n = 11$) for five

consecutive days. Frequencies of surface CCR1⁺ PBMC were significantly lower compared with PBS-treated BD mice ($n = 10$) ($4.59 \pm 1.36\%$ versus $7.06 \pm 1.8\%$, $P = 0.003$). The frequencies of surface CCR1⁺ LN cells were not different after CCL3 treatment compared with PBS control ($38.38 \pm 7.74\%$ versus $58.04 \pm 22.39\%$, $P = 0.01$). However, CCR1⁺ expression was not changed in PBMC cytoplasm (Fig. 4b). Treatment with CCL3 deteriorated BD symptoms in 10 (62.5%) of 16 BD mice and the disease severity score increased from 2.73 ± 0.47 to 3.2 ± 1.00 after 5 days of treatment. Four BD mice showed new skin ulcer and arthritis, and the remaining BD mice showed increased ulcer size.

BX471, a CCR1 antagonist, does not down-regulate CCR1⁺ cells in BD mice

BX471 was developed as a CCR1 antagonist to treat several inflammatory diseases, including rheumatoid




Figure 3. Frequencies of CCR1⁺ cells in CCL3-treated normal mice. CCL3 (50 pg and 100 pg) was intraperitoneally injected into normal mice for five consecutive days. Peripheral blood mononuclear cells (PBMC) and lymph node cells were then isolated. Frequencies of CCR1⁺ cells were analyzed by FACS on the surface (a) and in the cytoplasm (b) of PBMC and lymph node cells. Representative flow cytometric histograms of CCR1⁺ cells on the surface (a) and in the cytoplasm (b) of PBMC are shown in (c).

arthritis.¹² BX471 was administered to BD mice to down-regulate CCR1. BX471-treated BD mice ($n = 3$) showed $8.23 \pm 6.17\%$ of surface CCR1⁺ PBMC, $11.53 \pm 3.71\%$ of surface CCR1⁺ LN cells, $54.53 \pm 14.59\%$ of cytoplasm PBMC, and $67.17 \pm 11.49\%$ of cytoplasm LN cells. In BD mice, BX471 did not play a role in the down-regulation of CCR1⁺ cells (Fig. 4a,b). In normal mice, BX471-treated mice showed higher frequencies of surface CCR1⁺ PBMC than in PBS-treated normal mice ($17.22 \pm 8.66\%$ versus $12.15 \pm 7.92\%$, $P = 0.3$) and LN cells ($15.66 \pm 1.14\%$ versus $13.15 \pm 1.73\%$, $P = 0.05$). Inhibition of CCR1 expression by BX471 was only found in the cytoplasm of PBMC (BX471 versus PBS: $65.94 \pm 5.12\%$ versus $74.05 \pm 3.47\%$, $P = 0.02$) (Fig. 4c,d).

Drug treatment regulates frequencies of CCR1⁺ cells in normal mice

Colchicine and pentoxifylline are the most frequently prescribed medications for BD.²⁴ They are also used in BD mice.²⁵ To determine the correlation between CCR1 expression and these two agents, colchicine ($n = 7$) and pentoxifylline ($n = 7$) were administered to 4-week-old normal mice. The surface expression of CCR1 in PBMC was significantly up-regulated compared with PBS-treated normal mice ($n = 10$) (colchicine versus PBS: $16.02 \pm 2.2\%$ versus $11.11 \pm 2.39\%$, $P = 0.0006$; pentoxifylline versus PBS: $17.02 \pm 3.8\%$ versus $11.11 \pm 2.2\%$,

$P = 0.001$). Colchicine significantly up-regulated the expression of CCR1 on the surface of LN cells, compared with PBS-treated normal mice ($16.02 \pm 2.2\%$ versus $12.99 \pm 1.56\%$, $P = 0.005$). Pentoxifylline also up-regulated the expression of CCR1 compared with PBS-treated normal mice, although the up-regulation was not statistically significant ($14.8 \pm 3.6\%$ versus $12.99 \pm 1.56\%$, $P = 0.18$) (Fig. 5a). Colchicine and pentoxifylline significantly down-regulated the expression of cytoplasm CCR1⁺ PBMC ($59.37 \pm 11.5\%$, $P = 0.002$; $59.7 \pm 18.8\%$, $P = 0.03$, respectively) compared with PBS control ($73.65 \pm 3.0\%$). Cytoplasm CCR1 expression in LN cells was not altered under colchicine or pentoxifylline treatment (Fig. 5b).

Drug treatment regulates frequencies of CCR1⁺ cells in BD mice

Frequencies of CCR1⁺ cells were determined after BD mice were treated with medication. Treatment with colchicine ($n = 7$) or pentoxifylline ($n = 7$) significantly up-regulated CCR1 surface expression in PBMC of BD mice compared with PBS-treated BD mice ($n = 10$) ($15.92 \pm 4.5\%$ versus $7.06 \pm 1.8\%$, $P < 0.0001$; $19.083 \pm 3.4\%$ versus $7.06 \pm 1.8\%$, $P < 0.0001$, respectively). The frequencies of surface CCR1⁺ LN cells were also increased significantly by colchicine or pentoxifylline compared with the PBS-treated BD mice ($17.68 \pm 7.9\%$




Figure 4. Frequencies of CCR1⁺ cells in CCL3- and BX471-treated Behçet's disease (BD) mice. CCL3 and BX471 were used to treat BD mice and frequencies of CCR1⁺ cells were analyzed by FACS on the surface (a) and in the cytoplasm (b) of peripheral blood mononuclear cells (PBMC) and lymph node cells. Frequencies of CCR1⁺ cells in normal mice after treatment with BX471 are shown in (c) and (d).

versus $10.6 \pm 3.23\%$, $P = 0.02$; $21.36 \pm 4\%$ versus $10.6 \pm 6.9\%$, $P < 0.0001$, respectively) (Fig. 5c). In the cytoplasm of PBMC, treatment with colchicine and pentoxifylline increased the frequencies of CCR1⁺ cells. However, only pentoxifylline treatment resulted in a significant increase ($77.4 \pm 6.97\%$ versus $32.08 \pm 16.39\%$, $P < 0.0001$), colchicine or pentoxifylline did not affect the frequency of cytoplasm CCR1⁺ LN cells compared with PBS-treated BD mice (Fig. 5d). The histogram of CCR1⁺ expression in the surface and cytoplasm of PBMC after treatment with colchicine and pentoxifylline is presented in Fig. 5(e). Results show that the frequencies of surface CCR1⁺ expression rather than cytoplasm expression strongly correlated with the regulation of BD symptoms.

Anti-CCL3 antibody up-regulates the expression of CCR1⁺ cells and improves symptoms in BD mice

To determine whether CCR1 was regulated by CCL3 inhibition, anti-CCL3 antibody was intraperitoneally injected into normal mice. Frequencies of CCR1⁺ PBMC and LN cells were increased after anti-CCL3 antibody treatment. In surface staining of PBMC, treatment with anti-CCL3

antibody at 1–200 ng significantly increased frequencies of CCR1⁺ cells compared with PBS-treated normal mice (1 ng, $12.84 \pm 1.98\%$, $P = 0.009$; 2 ng, $13.37 \pm 2.28\%$, $P = 0.002$; 20 ng, $15.28 \pm 2.57\%$, $P < 0.0001$; 50 ng, $16.92 \pm 1.32\%$, $P < 0.0001$; 200 ng, $15.6 \pm 2.12\%$, $P = 0.0005$; versus PBS, $8.3 \pm 2.04\%$). In surface staining of LN cells, treatment with anti-CCL3 antibody significantly up-regulated frequencies of CCR1⁺ LN cells compared with PBS-treated normal mice (1 ng, $15.16 \pm 2.83\%$, $P < 0.0001$; 2 ng, $17.62 \pm 4.24\%$, $P < 0.0001$; 20 ng, $19.4 \pm 1.59\%$, $P < 0.0001$; 50 ng, $19.82 \pm 1.76\%$, $P < 0.0001$; 200 ng, $16.75 \pm 5.58\%$, $P = 0.0008$; versus PBS, $8.72 \pm 1.21\%$) (Fig. 6a). In the cytoplasm, frequencies of CCR1⁺ PBMC after treatment with anti-CCL3 antibody at 1–200 ng were not altered significantly in PBMC cytoplasm compared with PBS-treated normal mice (1 ng, $64.36 \pm 11.82\%$; 2 ng, $55.75 \pm 3.33\%$; 20 ng, $53.94 \pm 5.04\%$; 50 ng, $64.65 \pm 8.83\%$; 200 ng, $40.05 \pm 2.33\%$; versus PBS, $52.45 \pm 13.42\%$) (Fig. 6b). Cytoplasm CCR1⁺ LN cells were significantly up-regulated in mice treated with 1 ng of anti-CCL3 antibody ($42.12 \pm 8.93\%$, $P = 0.02$) or 200 ng of anti-CCL3 antibody ($59.0 \pm 7.77\%$, $P = 0.0002$) compared with PBS-treated normal mice




Figure 5. Frequencies of CCR1⁺ cells in colchicine (Col) and pentoxifylline (Pento) -treated normal and Behçet's disease (BD) mice. Colchicine (2 µg/mouse/day) and pentoxifylline (400 µg/mouse/day) were used to treat mice. Frequencies of surface and cytoplasm CCR1⁺ peripheral blood mononuclear cells (PBMC) and lymph node cells were analyzed in normal mice (a, b) and BD mice (c, d) by FACS. (e) Representative flow cytometric histogram of surface and cytoplasm CCR1⁺ PBMC following colchicine or pentoxifylline treatment of BD mice.




Figure 6. Frequencies of CCR1⁺ cells in anti-CCL3 antibody-treated normal mice. Anti-CCL3 antibody (1, 2, 20, 50 and 200 ng) was intraperitoneally injected into normal mice for five consecutive days. Peripheral blood mononuclear cells (PBMC) and lymph node cells were isolated. Frequencies of surface CCR1⁺ (a) and cytoplasm CCR1⁺ (b) PBMC and lymph node cells were analyzed by FACS.

(30.0 ± 7.27%). Treatment with 2, 20 and 50 ng of anti-CCL3 did not yield significant changes compared with PBS-treated normal mice (Fig. 6b).

To up-regulate CCR1⁺ cells in BD mice, anti-CCL3 antibody 20 ng/day was administered to BD mice ($n = 16$) for five consecutive days. The frequencies of CCR1⁺ expression on the surface of PBMC were significantly increased compared with those in PBS-treated BD mice ($n = 10$) (15.61 ± 2.92% versus 7.92 ± 2.61%, $P < 0.0001$). In LN cells, frequencies of surface CCR1⁺ expressions were significantly increased after anti-CCL3 antibody treatment (22.8 ± 2.7% versus 10.6 ± 3.1%, $P = 0.0001$) (Fig. 7a). In the cytoplasm of PBMC, the frequencies of CCR1 expression after treatment with anti-CCL3 antibody were up-regulated significantly compared with PBS control (51.42 ± 13.08% versus 32.11 ± 16.40%, $P < 0.0001$). However, the cytoplasm CCR1⁺ LN cells were down-regulated after treatment with anti-CCL3 antibody compared with PBS control (40.62 ± 7.81% versus 58.04 ± 22.43%, $P = 0.008$; Fig. 7b). Anti-CCL3 antibody treatment improved BD symptoms in 10 (50.0%) of 20 BD mice on day 5 after initiation with anti-CCL3 antibody treatment. The list of improved mice is provided in the Supplementary material (Table S1). The disease severity score in BD mice changed from 3.1 ± 0.3 to 2.45 ± 0.68 on day 5 with symptom improvement after anti-CCL3 antibody treatment

($P < 0.0001$). Disease severity score was also significantly varied between CCL3-treated BD mice (3.2 ± 1.0) and anti-CCL3 antibody-treated BD mice (2.45 ± 0.68) on day five ($P = 0.004$) (Fig. 7c). Figure 7(d) show the changes in symptoms after treatment with CCL3 or anti-CCL3 antibody in BD mice.

Plasma levels of CCL3 in patients with BD

The levels of CCL3 in patients with active BD (76.88 ± 74.48 pg/ml) were higher than in patients with inactive BD (24.58 ± 7.54 pg/ml, $P = 0.09$) and healthy control individuals (22.82 ± 7.87 pg/ml, $P = 0.02$) (Fig. 8).

Discussion

This study quantitatively analyzed the expression of chemokine receptors in PBMC and LN cells of BD mice. Our results suggest that BD-like symptoms were correlated with frequencies of CCR1⁺ cells. Percentages of CCR1⁺ cells were significantly down-regulated in BD mice compared with normal or BDN mice. Cytokine IL-10- and GM-CSF-treated BD mice exhibited significant up-regulation of CCR1⁺ cells compared with PBS-treated BD mice. CCR1 ligand CCL3-treated BD mice showed significant down-regulation of CCR1⁺ cells compared with PBS-




Figure 7. Frequencies of CCR1⁺ cells in anti-CCL3 antibody-treated Behçet's disease (BD) mice. Anti-CCL3 antibody 20 ng was intraperitoneally injected into BD mice for five consecutive days. Peripheral blood mononuclear cells (PBMC) and lymph node cells were then isolated. Frequencies of surface CCR1⁺ (a) and cytoplasm CCR1⁺ (b) PBMC and lymph node cells were analyzed by FACS. (c) The changes in disease severity score after CCL3 or anti-CCL3 antibody treatment of BD mice. (d) Symptoms of BD mice after CCL3 or anti-CCL3 antibody treatment.




Figure 8. Plasma levels of CCL3 in patients with active and inactive Behçet's disease (BD).

treated BD mice. Ten of 16 CCL3-treated BD mice showed deteriorated BD symptoms. Anti-CCL3 antibody treatment improved BD symptoms, decreased the disease severity score, and up-regulated CCR1⁺ cells significantly compared with PBS-treated BD mice. Treatment of BD mice with colchicine and pentoxifylline significantly up-

regulated CCR1⁺ cells compared with PBS treatment. Frequencies of CCR1⁺ cells were strongly associated with symptoms in BD mice.

Chemokine receptors are essential moderators of leukocyte trafficking in inflammatory diseases. CCR1 may be a strong therapeutic target in a variety of diseases.²⁶ CCR1 has been shown to play a role in recruiting monocytes and Th1 cells according to active inflammation.²⁷ CCR1 and its ligands are associated with the development of experimental arthritis. In rheumatoid arthritis, CCR1 is an efficient therapeutic target because it mediates the regulation of tumor necrosis factor- α and IL-10.²⁸ However, in patients with BD, the frequencies of CCR1⁺ cells are decreased in patients with active BD than in normal controls, even though the difference was not statistically significant.²⁹ Similarly, this study also shows that the frequencies of CCR1⁺ cells in BD mice were down-regulated compared with the control mice. Down-regulated CCR1 expression has been reported after infection with *Leishmania infantum*³⁰ or coronavirus.³¹ Correlation between HSV infection and CCR1 expression does not show consistent results.^{32,33}

Chemokine CCL3 expression has been reported in arthritic inflammation.³⁴ CCL3 recruits inflammatory leukocytes by binding to CCR1.³⁵ It affects chronic joint inflammation.³⁶ Neutralizing antibodies to CCL3 alleviate inflammation.³⁷ Higher serum levels of CCL3 have been found in patients with active BD than in patients with inactive stages of BD.³⁸ Results of the present study demonstrate that treatment with CCL3 deteriorated symptoms in BD mice and down-regulated the frequencies of CCR1⁺ cells. Increased CCL3 expression exacerbates BD symptoms through down-regulation of CCR1.

Interleukin-10 is known as a CCR1 stimulator.³⁹ It is mainly expressed in monocytes, macrophages, Th2 cells, and B cells. IL-10 is one of the key anti-inflammatory cytokines expressed in the human immune response. It is a strong inhibitor of Th1 cytokines IL-2 and interferon- γ .^{40,41} Anti-inflammatory cytokine IL-10 suppresses pro-inflammatory cytokines and protects the host against tissue damage.⁴² In patients with BD, IL-10 polymorphisms have been identified.^{43,44} Such gene variation might be correlated with insufficient expression of IL-10. Results of the present study show that IL-10 up-regulated the frequencies of CCR1⁺ cells in BD mice compared with PBS control. IL-10 and GM-CSF also synergistically increased the frequencies of CCR1⁺ cells in BD mice.

Granulocyte-macrophage colony-stimulating factor is a stimulator of CCR1 expression.²² It enhances the maturation of dendritic cells and plays a role in recruiting circulating neutrophils, monocytes and lymphocytes for host defense.⁴⁵ Different types of inflammatory and immune cells such as dendritic cells, monocytes, neutrophils, microglia and macrophages are modulated by GM-CSF under autoimmune conditions.⁴⁶ GM-CSF may influence chronic inflammatory diseases such as arthritis,⁴⁷ experimental autoimmune encephalitis⁴⁸ and airway inflammation.⁴⁹ GM-CSF also enhances resistance to viral infection⁵⁰ and bacterial translocation.⁵¹ In patients with BD, GM-CSF transcription was down-regulated more than in controls.⁵² However, the protein expression or serum levels of GM-CSF in patients with BD have yet to be elucidated. According to a case report, intra-lesional injection with recombinant human GM-CSF induces wound healing in genital ulcers of BD.⁵³ GM-CSF also enhances CCR1 expression.^{21,22} Our data demonstrated that treatment with GM-CSF significantly up-regulated CCR1⁺ expression in PBMC and LN cells.

BX471 was developed as a potent antagonist against CCR1.⁵⁴ BX471 treatment attenuated systemic inflammation of sepsis,⁵⁵ nephritis,⁵⁶ arthritis⁵⁷ and myocarditis.⁵⁸ In our study, BX471 was administered to BD mice to confirm the correlation of CCR1 expression with BD by inhibiting CCR1. BX471 treatment failed to inhibit the frequencies of surface or cytoplasm CCR1⁺ PBMC and LN cells in BD mice. In normal mice, BX471 inhibited the expression of CCR1 only in the cytoplasm of PBMC.

These results indicate that BX471 treatment failed to adequately inhibit CCR1⁺ cells *in vivo*, which partly explains why clinical trials of BX471 inhibition of CCR1 were discontinued.

In conclusion, our study demonstrated down-regulated frequencies of CCR1⁺ cells, which were closely correlated with BD symptoms and the expression of CCR1⁺ cells was controlled by cytokine IL-10, GM-CSF and ligand CCL3. Even though BD is a multi-factorial inflammatory disease associated with immune dysregulation, treatment with single chemokine or chemokine receptor alone contributes to immune regulation resulting in the control of systemic symptoms of BD.

Acknowledgements

SS designed experiments and analyzed the data. HMS and HOB performed experiments. ESL supplied and was consulted on human samples. HMS and SS wrote the manuscript. This study was supported by a grant from the Korea Health Technology R&D Project through the Korea Health Industry Development Institute funded by the Ministry of Health & Welfare, Republic of Korea (HI15C2483) and the Basic Science Research Program through the National Research Foundation of Korea funded by the Ministry of Education (2017R1D1A1B03032168).

Disclosure

The authors declare no conflicts of interest.

References

- Hou S, Xiao X, Li F, Jiang Z, Kijlstra A, Yang P. Two-stage association study in Chinese Han identifies two independent associations in CCR1/CCR1 locus as candidate for Behçet's disease susceptibility. *Hum Genet* 2012; **131**:1841–50.
- Kirino Y, Bertias G, Ishigatsubo Y, Mizuki N, Tugal-Tutkun I, Seyahi E *et al.* Genome-wide association analysis identifies new susceptibility loci for Behçet's disease and epistasis between HLA-B*51 and ERAP1. *Nat Genet* 2013; **45**:202–7.
- Su SB, Mukaida N, Wang J, Nomura H, Matsushima K. Preparation of specific polyclonal antibodies to a C-C chemokine receptor, CCR3, and determination of CCR3 expression on various types of leukocytes. *J Leukoc Biol* 1996; **60**:658–66.
- Cochand L, Isler P, Songeon F, Nicod LP. Human lung dendritic cells have an immature phenotype with efficient mannose receptors. *Am J Respir Cell Mol Biol* 1999; **21**:547–54.
- Gao JL, Wynn TA, Chang Y, Lee EJ, Broxmeyer HE, Cooper S *et al.* Impaired host defense, hematopoiesis, granulomatous inflammation and type 1-type 2 cytokine balance in mice lacking CC chemokine receptor 1. *J Exp Med* 1997; **185**:1959–68.
- Bonecchi R, Polentarutti N, Luini W, Borsatti A, Bernasconi S, Locati M *et al.* Up-regulation of CCR6 and CCR6 and induction of chemotaxis to CC chemokines by IFN- γ in human neutrophils. *J Immunol* 1999; **162**:474–9.
- Katschke KJ Jr, Rottman JB, Ruth JH, Qin S, Wu L, LaRosa G *et al.* Differential expression of chemokine receptors on peripheral blood, synovial fluid, and synovial tissue monocytes/macrophages in rheumatoid arthritis. *Arthritis Rheum* 2001; **44**:1022–32.
- Joubert P, Lajoie-Kadoch S, Welman M, Dragon S, Letuvee S, Tolloczko B *et al.* Expression and regulation of CCR8 by airway smooth muscle cells in asthma. *J Immunol* 2008; **180**:1268–75.
- Bignon A, Gaudin F, Hemon P, Tharinger H, Mayol K, Walzer T *et al.* CCR9 inhibition ameliorates the progression of lupus nephritis in NZB/W mice. *J Immunol* 2014; **192**:886–96.

- 10 Blease K, Mehrad B, Standiford TJ, Lukacs NW, Kunkel SL, Chensue SW *et al.* Airway remodeling is absent in CCR10^{-/-} mice during chronic fungal allergic airway disease. *J Immunol* 2000; **165**:1564–72.
- 11 Choi SW, Hildebrandt GC, Oikiewicz KM, Hanauer DA, Chaudhary MN, Silva IA *et al.* CCR11/CCL5 (RANTES) receptor–ligand interactions modulate allogeneic T-cell responses and graft-versus-host disease following stem-cell transplantation. *Blood* 2007; **110**:3447–55.
- 12 Horuk R. Chemokine receptor antagonists: overcoming developmental hurdles. *Nat Rev Drug Discov* 2009; **8**:23–33.
- 13 Eglin RP, Lehner T, Subak-Sharpe JH. Detection of RNA complementary to herpes-simplex virus in mononuclear cells from patients with Behçet's syndrome and recurrent oral ulcers. *Lancet* 1982; **2**:1356–61.
- 14 Young C, Lehner T, Barnes CG. CD4 and CD8 cell responses to herpes simplex virus in Behçet's disease. *Clin Exp Immunol* 1988; **73**:6–10.
- 15 Behcet H. Über rezidivierende, aphthöse, durch ein virus verursachte geschwüre am mund, am auge und an den genitalien. *Dermatologische Wochenschrift* 1937; **105**:1152–7.
- 16 Sohn S, Lee ES, Bang D, Lee S. Behçet's disease-like symptoms induced by the Herpes simplex virus in ICR mice. *Eur J Dermatol* 1998; **8**:21–3.
- 17 Sohn S, Lee ES, Bang D. Learning from HSV-infected mice as a model of Behçet's disease. *Clin Exp Rheumatol* 2012; **30**:596–103.
- 18 Kim DY, Cho S, Choi MJ, Sohn S, Lee ES, Bang D. Immunopathogenic role of herpes simplex virus in Behçet's disease. *Genet Res Int* 2013; **2013**:638273.
- 19 Zheng Z, Sohn S, Ahn KJ, Bang D, Cho SB. Serum reactivity against herpes simplex virus type 1 UL48 protein in Behçet's disease patients and a Behçet's disease-like mouse model. *Acta Derm Venereol* 2015; **95**:952–8.
- 20 Vaiopoulos GKP, Evangelatos N, Kaklamanis PH. The spectrum of mucocutaneous manifestations in Adamantiades-Behçet's disease in Greece. *J Eur Acad Dermatol Venerol* 2010.
- 21 Li H, Cheung W, Choo HH, Chan JH, Lai PS, Fred Wong WS. IL-10 synergistically enhances GM-CSF-induced CCR21 expression in myelomonocytic cells. *Biochem Biophys Res Commun* 2003; **304**:417–24.
- 22 Cheng SS, Lai JJ, Lukacs NW, Kunkel SL. Granulocyte–macrophage colony stimulating factor up-regulates CCR22 in human neutrophils. *J Immunol* 2001; **166**:1178–84.
- 23 Elsner J, Dulkys Y, Gupta S, Escher SE, Forssmann WG, Kapp A *et al.* Differential pattern of CCR23 internalization in human eosinophils: prolonged internalization by CCL5 in contrast to CCL3. *Allergy* 2005; **60**:1386–93.
- 24 Bang D. Treatment of Behçet's disease. *Yonsei Med J* 1997; **38**:401–10.
- 25 Choi B, Kim HA, Suh CH, Byun HO, Jung JY, Sohn S. The relevance of miRNA-21 in HSV-induced inflammation in a mouse model. *Int J Mol Sci* 2015; **16**:7413–27.
- 26 Gerard C, Rollins BJ. Chemokines and disease. *Nat Immunol* 2001; **2**:108–15.
- 27 Oo YH, Shetty S, Adams DH. The role of chemokines in the recruitment of lymphocytes to the liver. *Dig Dis* 2010; **28**:31–44.
- 28 Amat M, Benjamim CF, Williams LM, Prats N, Terracabras E, Beleta J *et al.* Pharmacological blockade of CCR28 ameliorates murine arthritis and alters cytokine networks *in vivo*. *Br J Pharmacol* 2006; **149**:666–75.
- 29 Houman H, Hamzaoui A, Ben Ghorbal I, Khanfir M, Feki M, Hamzaoui K. Abnormal expression of chemokine receptors in Behçet's disease: relationship to intracellular Th1/Th2 cytokines and to clinical manifestations. *J Autoimmun* 2004; **23**:267–73.
- 30 Panaro MA, Spinelli R, Lisi S, Sisto M, Acquafredda A, Fumarola L *et al.* Reduced expression of the chemokine receptor CCR30 in human macrophages and U-937 cells *in vitro* infected with *Leishmania infantum*. *Clin Exp Med* 2004; **3**:225–30.
- 31 Hickey MJ, Held KS, Baum E, Gao JL, Murphy PM, Lane TE. CCR31 deficiency increases susceptibility to fatal coronavirus infection of the central nervous system. *Viral Immunol* 2007; **20**:599–608.
- 32 Cook WJ, Kramer MF, Walker RM, Burwell TJ, Holman HA, Coen DM *et al.* Persistent expression of chemokine and chemokine receptor RNAs at primary and latent sites of herpes simplex virus 1 infection. *Virology* 2004; **1**:5.
- 33 Sorensen LN, Paludan SR. Blocking CC chemokine receptor (CCR) 1 and CCR3 during herpes simplex virus type 2 infection *in vivo* impairs host defence and perturbs the cytokine response. *Scand J Immunol* 2004; **59**:321–33.
- 34 Zhang R, Sun P, Jiang Y, Chen Z, Huang C, Zhang X *et al.* Genome-wide haplotype association analysis and gene prioritization identify CCL3 as a risk locus for rheumatoid arthritis. *Int J Immunogenet* 2010; **37**:273–8.
- 35 Turner MD, Nedjai B, Hurst T, Pennington DJ. Cytokines and chemokines: at the crossroads of cell signalling and inflammatory disease. *Biochim Biophys Acta* 2014; **1843**:2563–82.
- 36 Chintalacharuvu SR, Wang JX, Giaconia JM, Venkataraman C. An essential role for CCL3 in the development of collagen antibody-induced arthritis. *Immunol Lett* 2005; **100**:202–4.
- 37 Manczak M, Jiang S, Orzechowska B, Adamus G. Crucial role of CCL3/MIP-1 α in the recurrence of autoimmune anterior uveitis induced with myelin basic protein in Lewis rats. *J Autoimmun* 2002; **18**:259–70.
- 38 Kim WU, Do JH, Park KS, Cho ML, Park SH, Cho CS *et al.* Enhanced production of macrophage inhibitory protein-1 α in patients with Behçet's disease. *Scand J Rheumatol* 2005; **34**:129–35.
- 39 Musso T, Cappello P, Stornello S, Ravarino D, Caorsi C, Otero K *et al.* IL-10 enhances CCL2 release and chemotaxis induced by CCL16 in human monocytes. *Int J Immunopathol Pharmacol* 2005; **18**:339–49.
- 40 Li L, Elliott JF, Mosmann TR. IL-10 inhibits cytokine production, vascular leakage, and swelling during T helper 1 cell-induced delayed-type hypersensitivity. *J Immunol* 1994; **153**:3967–78.
- 41 Saraiva M, O'Garra A. The regulation of IL-10 production by immune cells. *Nat Rev Immunol* 2010; **10**:170–81.
- 42 Cytkot JC, Turner J. Interleukin-10 and immunity against prokaryotic and eukaryotic intracellular pathogens. *Infect Immun* 2011; **79**:2964–73.
- 43 Mizuki N, Meguro A, Ota M, Ohno S, Shiota T, Kawagoe T *et al.* Genome-wide association studies identify IL23R-IL12RB2 and IL10 as Behçet's disease susceptibility loci. *Nat Genet* 2010; **42**:703–6.
- 44 Rimmers EF, Cosan F, Kirino Y, Ombrello MJ, Abaci N, Satorius C *et al.* Genome-wide association study identifies variants in the MHC class I, IL10, and IL23R-IL12RB2 regions associated with Behçet's disease. *Nat Genet* 2010; **42**:698–702.
- 45 Shi Y, Liu CH, Roberts AI, Das J, Xu G, Ren G *et al.* Granulocyte–macrophage colony-stimulating factor (GM-CSF) and T-cell responses: what we do and don't know. *Cell Res* 2006; **16**:126–33.
- 46 Bhattacharya P, Thirupathi M, Elshabrawy HA, Alharshawi K, Kumar P, Prabhakar BS. GM-CSF: an immune modulatory cytokine that can suppress autoimmunity. *Cytokine* 2015; **75**:261–71.
- 47 Cornish AL, Campbell IK, McKenzie BS, Chatfield S, Wicks IP. G-CSF and GM-CSF as therapeutic targets in rheumatoid arthritis. *Nat Rev Rheumatol* 2009; **5**:554–9.
- 48 Ponomarev ED, Shriver LP, Maresz K, Pedras-Vasconcelos J, Verthelyi D, Dittel BN. GM-CSF production by autoreactive T cells is required for the activation of microglial cells and the onset of experimental autoimmune encephalomyelitis. *J Immunol* 2007; **178**:39–48.
- 49 Cates EC, Fattouh R, Wattie J, Inman MD, Goncharova S, Coyle AJ *et al.* Intranasal exposure of mice to house dust mite elicits allergic airway inflammation via a GM-CSF-mediated mechanism. *J Immunol* 2004; **173**:6384–92.
- 50 Huang FF, Barnes PF, Feng Y, Donis R, Chronoes ZC, Idell S *et al.* GM-CSF in the lung protects against lethal influenza infection. *Am J Respir Crit Care Med* 2011; **184**:259–68.
- 51 Unal AE, Cevikel MH, Ozgun H, Tunger A. Effect of granulocyte–macrophage colony stimulating factor on bacterial translocation after experimental obstructive jaundice. *Eur J Surg* 2001; **167**:366–70.
- 52 Takahama H, Itoh R, Inoue-Komatsu C, Furusawa S, Takahashi H, Mizoguchi M. Granulocyte colony-stimulating factor (G-CSF) and granulocyte–macrophage colony-stimulating factor (GM-CSF) in Behçet's disease. *J Dermatol* 1994; **21**:546–52.
- 53 Alli N, Karakayali G, Kahraman I, Artuz F. Local intralesional therapy with rhGM-CSF for a large genital ulcer in Behçet's disease. *Br J Dermatol* 1997; **136**:639–40.
- 54 Vaidehi N, Schlyer S, Trabanino RJ, Floriano WB, Abrol R, Sharma S *et al.* Predictions of CCR54 chemokine receptor structure and BX 471 antagonist binding followed by experimental validation. *J Biol Chem* 2006; **281**:27613–20.
- 55 He M, Horuk R, Mochhala SM, Bhatia M. Treatment with BX471, a CC chemokine receptor 1 antagonist, attenuates systemic inflammatory response during sepsis. *Am J Physiol Gastrointest Liver Physiol* 2007; **292**:G1173–80.
- 56 Anders HJ, Belemzova E, Eis V, Segerer S, Vielhauer V, Perez de Lema G *et al.* Late onset of treatment with a chemokine receptor CCR56 antagonist prevents progression of lupus nephritis in MRL-Fas(lpr) mice. *J Am Soc Nephrol* 2004; **15**:1504–13.
- 57 Lebre MC, Vergunst CE, Choi IY, Aarras S, Oliveira AS, Wyant T *et al.* Why CCR57 and CCR57 blockade failed and why CCR57 blockade might still be effective in the treatment of rheumatoid arthritis. *PLoS ONE* 2011; **6**:e21772.
- 58 Futamatsu H, Suzuki J, Koga N, Adachi S, Kosuge H, Maejima Y *et al.* A CCR58 antagonist prevents the development of experimental autoimmune myocarditis in association with T cell inactivation. *J Mol Cell Cardiol* 2006; **40**:853–61.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1 List of symptoms in anti-CCL3 antibody-treated Behçet's disease mice.